

## Author Index

- Abdu, T.A.M. *see* Barton, D.M.  
Abe, T. *see* Nagai, Y.  
Acerini, C.L. *see* Ford-Adams, M.E.  
Adamson, U. *see* Bragg, J.  
Adlis, S.A. *see* Mazze, R.S.  
Adriaanse, M.C., Snoek, F.J., Dekker, J.M., Spijkerman, A.M.W., Nijpels, G., van der Ploeg, H.M. & Heine, R.J. Perceived risk for Type 2 diabetes in participants in a stepwise population-screening programme, 210  
Agrawal, A. & McKibbin, M.A. Technical failure in photographic screening for diabetic retinopathy, 777  
Ahn, C.W. *see* Kim, D.J.  
Aitken, G. *see* Sommerfield, A.J.  
Al-Attas, O. *see* Al-Daghri, N.  
Al-Daghri, N., Al-Rubean, K., Bartlett, W.A., Al-Attas, O., Jones, A.F. & Kumar, S. Serum leptin is elevated in Saudi Arabian patients with metabolic syndrome and coronary artery disease, 832  
Al-Rubean, K. *see* Al-Daghri, N.  
Alazri, M.H. & Neal, R.D. The association between satisfaction with services provided in primary care and outcomes in Type 2 diabetes mellitus, 486  
Albert, A. *see* Zhang, L.  
Alberti, K.G.M.M. *see* Fischbacher, C.M.  
Alberti, K.G.M.M. *see* Parker, L.  
Alberti, K.G.M.M. *see* Williams, J.W.  
Alberti, K.G.M.M. *see* Zimmer, P.  
Albrecht, M. *see* Ortlepp, J.R.  
Alcolado, J. *see* Channon, S.  
Aldington, S. *see* Harding, S.  
Aldington, S.J. *see* Scanlon, P.H.  
Alhenc-Gelas, F. *see* Hadjadj, S.  
Allen, K.V. *see* Henderson, J.N.  
Almind, K. *see* Pildal, J.  
Álvarez, L. *see* Costa, A.  
Aman, J. *see* Ingberg, C.M.  
Amano, S. *see* Pang, H.  
Ambye, L. *see* Pildal, J.  
Amiel, S.A. *see* Bingham, E.  
Amin, P. & Sturrock, N.D.C. A pilot study of the beneficial effects of amantadine in the treatment of painful diabetic peripheral neuropathy, 114  
Amod, A. *see* Motala, A.A.  
Anderson, B. *see* Welch, G.  
Anderson, J. *see* Barton, D.M.  
Anderson, P. *see* Lunt, H.  
Anderson, R.L. *see* Mazze, R.S.  
Andreasen, A.H. *see* Nordly, S.  
Anshoo, H. *see* Velmurugan, K.  
Aoki, T. *see* Yokoyama, H.  
Araújo-Vilar, D. *see* García-Estévez, D.A.  
Armstrong, D.G. & Frykberg, R.G. Classifying diabetic foot surgery: toward a rational definition, 329  
Armstrong, D.G., Dang, C., Nixon, B.P. & Boulton, A.J.M. The hazards of the holiday foot: persons at high risk for diabetic foot ulceration may be more active on holiday, 247  
Arun, C.S., Ngugi, N., Lovelock, L. & Taylor, R. Effectiveness of screening in preventing blindness due to diabetic retinopathy, 186  
Aso, Y., Okumura, K., Yoshida, N., Tayama, K., Kanda, T., Kobayashi, I., Takemura, Y. & Inukai, T. Plasma interleukin-6 is associated with coagulation in poorly controlled patients with Type 2 diabetes, 930  
Bachmann, M.O., Eachus, J., Hopper, C.D., Davey Smith, G., Propper, C., Pearson, N.J., Williams, S., Tallon, D. & Frankel, S. Socio-economic inequalities in diabetes complications, control, attitudes and health service use: a cross-sectional study, 921  
Baik, S.H. *see* Choi, K.M.  
Bain, S.C., Gill, G.V., Dyer, P.H., Jones, A.F., Murphy, M., Jones, K.E., Smyth, C. & Barnett, A.H. Characteristics of Type 1 diabetes of over 50 years duration (the Golden Years Cohort), 808  
Baldeweg, S.E. & Rosenthal, M. The 20th Annual Meeting of the Anglo-Danish-Dutch Diabetes Group, 332  
Banerjee, K. *see* Saeed, B.O.  
Bardelli, C. *see* Bo, S.  
Bardini, G. *see* Mannucci, E.  
Barnett, A.H. A review of basal insulins, 873  
Barnett, A.H. *see* Bain, S.C.  
Barnett, A.H., Grant, P.J., Hitman, G.A., Mather, H., Pawa, M., Robertson, L. & Trelfa, A. on behalf of the Indo-Asian Trial Investigators. Rosiglitazone in Type 2 diabetes mellitus: an evaluation in British Indo-Asian patients, 387  
Bartlett, W.A. *see* Al-Daghri, N.  
Barton, D.M., Baskar, V., Kamalakannan, D., Buch, H.N., Gone, K., Wilson, E., Anderson, J. & Abdu, T.A.M. An assessment of care of paediatric and adolescent patients with diabetes in a large district general hospital, 394  
Baskar, V. *see* Barton, D.M.  
Basu, A., Kamal, A.D., Illahi, W., Khan, M., Stavrou, P. & Ryder, R.E.J. Is digital image compression acceptable within diabetic retinopathy screening?, 766  
Baxter, M. *see* Masson, E.A.  
Beck-Nielsen, H. *see* Jensen, D.M.  
Beck-Nielsen, H. *see* Staehr, P.  
Becker, D. *see* Bosnyak, Z.  
Bek, T. *see* Frystyk, J.  
Ben Mehidi, A. *see* Massin, P.  
Ben-Haroush, A. *see* Yogeve, Y.  
Bengel, F.M. *see* Hattori, N.  
Bennett, S.M.A. *see* Parker, L.  
Bergental, R.M. *see* Mazze, R.S.  
Berghold, A. *see* Siebenhofer, A.

- Berglund, G. *see* Gottsäter, A.  
 Bergrem, H. *see* Schramm, W.  
 Berne, C. *see* Ingberg, C.-M.  
 Bhattarai, M.D. *see* Singh, D.L.  
 Bhopal, R. *see* Fischbacher, C.M.  
 Biermann, E. No evidence for accumulation of insulin glargine (LANTUS), 333  
 Biesenbach, G., Raml, A., Schmekal, B. & Eichbauer-Sturm, G. Decreased insulin requirement in relation to GFR in nephropathic Type 1 and insulin-treated Type 2 diabetic patients, 642  
 Bingham, E., Hopkins, D., Pernet, A., Reid, H., Macdonald, I.A. & Amiel, S.A. The effects of KATP channel modulators on counterregulatory responses and cognitive function during acute controlled hypoglycaemia in healthy men: a pilot study, 231  
 Birkeland, K.I., Kilhovd, B., Thorsby, P., Torjesen, P.A., Ganss, R., Vaaler, S. & Hanssen, K.F. Heterogeneity of non-insulin-dependent diabetes expressed as variability in insulin sensitivity,  $\beta$ -cell function and cardiovascular risk profile, 37  
 Biswas, M. *see* Prakash, P.K.  
 Bjerle, P. *see* Lindmark, S.  
 Blaak, E.E. *see* Mensink, M.  
 Bo, S., Menato, G., Pinach, S., Signorile, A., Bardelli, C., Lezo, A., Marchisio, B., Gentile, L., Cassader, M., Massobrio, M. & Pagano, G. Clinical characteristics and outcome of pregnancy in women with gestational hyperglycaemia with and without antibodies to  $\beta$ -cell antigens, 64  
 Bodansky, H.J. *see* Feltbower, R.G.  
 Bodansky, H.J. *see* McKinney, P.A.  
 Bolli, G. *see* Owens, D.R.  
 Borch-Johnsen, K. *see* Colagiuri, S.  
 Borch-Johnsen, K. *see* Pildal, J.  
 Borch-Johnsen, K. *see* Schramm, W.  
 Borch-Johnsen, K., Lauritzen, T., Glümer, C. & Sandbæk, A. Screening for Type 2 diabetes—should it be now? 175  
 Bos, G. *see* Scheffer, P.G.  
 Bosnyak, Z., Forrest, K.Y.-Z., Maser, R.E., Becker, D. & Orchard, T.J. Do plasminogen activator inhibitor (PAI-1) or tissue plasminogen activator PAI-1 complexes predict complications in Type 1 diabetes: The Pittsburgh Epidemiology of Diabetes Complications Study, 147  
 Botas, P., Delgado, E., Castaño, G., Díaz de Greñu, C., Prieto, J. & Díaz-Cadorniga, F.J. Comparison of the diagnostic criteria for diabetes mellitus, WHO-1985, ADA-1997 and WHO-1999 in the adult population of Asturias (Spain), 904  
 Bott, U., Ebrahim, S., Hirschberger, S. & Skovlund, S.E. Effect of the rapid-acting insulin analogue insulin aspart on quality of life and treatment satisfaction in patients with Type 1 diabetes, 626  
 Boucher, B.J. Trends in the incidence of childhood diabetes in South Asians and other children in Bradford, UK, 602  
 Boulton, A.J.M. *see* Armstrong, D.G.  
 Boulton, A.J.M. *see* Dang, C.N.  
 Bowers, A. *see* Manning, P.J.  
 Boyce, K.S. *see* Mazze, R.S.  
 Bragd, J., Adamson, U., Lins, P.-E., Wredling, R. & Oskarsson, P. A repeated cross-sectional survey of severe hypoglycaemia in 178 Type 1 diabetes mellitus patients performed in 1984 and 1998, 216  
 Braithwaite, P.A. *see* Prakash, P.K.  
 Brash, P.D. *see* Masson, E.A.  
 Brauer, M. *see* Schaefer-Graf, U.M.  
 Briggs, A. *see* Clarke, P.  
 Broadbent, D.M. *see* Younis, N.  
 Brown, K. *see* Saeed, B.O.  
 Bruce, D.G. *see* Davis, T.M.E.  
 de Bruin, T.W.A. *see* Mensink, M.  
 Brydon, P. *see* Dunne, F.  
 Brydon, P. *see* Sinha, B.  
 Buch, H.N. *see* Barton, D.M.  
 Buchanan, T.A. *see* Chong-Martinez, B.  
 Bühling, K.J. *see* Schaefer-Graf, U.M.  
 Bulsara, M. *see* Caplin, N.J.  
 Bury, G. *see* Smith, S.M.  
 Busch, K. & Chantelau, E. Effectiveness of a new brand of stock 'diabetic' shoes to protect against diabetic foot ulcer relapse. A prospective cohort study, 665  
 Buxton-Thomas, M.S. *see* Watkins, P.J.  
 Byrne, K. *see* McFarlane, S.I.  
 Cabezas-Cerrato, J. *see* García-Estévez, D.A.  
 Caldwell, G. *see* Masson, E.A.  
 Cameron, A.J. *see* Williams, J.W.  
 Cameron, F.J., Smids, D., Hesketh, K., Wake, M. & Northam, E.A. Early detection of emotional and behavioural problems in children with diabetes: the validity of the Child Health Questionnaire as a screening instrument, 646  
 Campbell, L.V. *see* Kriketos, A.D.  
 Caplin, N.J., O'Leary, P., Bulsara, M., Davis, E.A. & Jones, T.W. Subcutaneous glucose sensor values closely parallel blood glucose during insulin-induced hypoglycaemia, 238  
 Carey, D.G. *see* Kriketos, A.D.  
 Carr, L. *see* Scuffham, P.  
 Carroll, C. *see* Dornan, T.  
 Carroll, C., Naylor, E., Marsden, P. & Dornan, T. How do people with Type 2 diabetes perceive and respond to cardiovascular risk? 355  
 Carstensen, B. *see* Colagiuri, S.  
 Carstensen, B. *see* Pildal, J.  
 Casals, E. *see* Costa, A.  
 Casamitjana, R. *see* Costa, A.  
 Cassader, M. *see* Bo, S.  
 Castaño, G. *see* Botas, P.  
 Catto, A.J. *see* Kain, K.  
 Cerimele, B. *see* Roach, P.  
 Cha, B.S. *see* Kim, D.J.  
 Chan, J.C.N. *see* Ko, G.T.C.  
 Chan, J.C.N. *see* Thomas, G.N.  
 Chan, W.B. *see* Ko, G.T.C.  
 Chang, H.-Y. *see* Pan, W.-H.  
 Channon, S., Smith, V., Alcolado, J. & Gregory, J.W. Current methods of transfer of young people with Type 1 diabetes to adult services, 1034  
 Chantelau, E. *see* Busch, K.  
 Chatellier, G. *see* Hadjadj, S.  
 Chatterjee, S., Tsaloumas, M.D., Gee, H., Lipkin, G. & Dunne, F.P. From minimal background diabetic retinopathy to profuse sight threatening vitreoretinal haemorrhage: management issues in a case of pregestational diabetes and pregnancy, 683  
 Chen, R. *see* Yoge, Y.  
 Chen, T.H.-H. *see* Liu, W.-J.  
 Chisholm, D.J. *see* Kriketos, A.D.

- Chitson, P. *see* Williams, J.W.
- Chiu, K.C., Chuang, L.-M., Chu, A. & Wang, M. Transcription factor 1 and  $\beta$ -cell function in glucose-tolerant subjects, 225
- Choi, D.S. *see* Choi, K.M.
- Choi, K.M., Lee, J., Kim, K.B., Kim, D.R., Kim, S.K., Shin, D.H., Kim, N.H., Park, I.B., Choi, D.S. & Baik, S.H. Factor analysis of the metabolic syndrome among elderly Koreans—the South-west Seoul Study, 99
- Chong-Martinez, B., Buchanan, T.A., Wenby, R.B. & Meiselman, H.J. Decreased red blood cell aggregation subsequent to improved glycaemic control in Type 2 diabetes mellitus, 301
- Christensen, J. *see* Colagiuri, S.
- Christie, D. *see* Viner, R.M.
- Chu, A. *see* Chiu, K.C.
- Chuang, L.-M. *see* Chiu, K.C.
- Citroni, N. *see* Inchiostro, S.
- Clark, M. & Hampson, S.E. Comparison of patients' and healthcare professionals' beliefs about and attitudes towards Type 2 diabetes, 152
- Clarke, P., Gray, A., Legood, R., Briggs, A. & Holman, R. The impact of diabetes-related complications on healthcare costs: results from the United Kingdom Prospective Diabetes Study (UKPDS Study No. 65), 442
- Clarke, P. *see* Holt, R.I.G.
- Cockram, C.S. *see* Ko, G.T.C.
- Cockram, C.S. *see* Thomas, G.N.
- Colagiuri, S., Sandbæk, A., Carstensen, B., Christensen, J., Glumer, C., Lauritzen, T. & Borch-Johnsen, K. Comparability of venous and capillary glucose measurements in blood, 953
- Coleman, M.A.G. *see* Holt, R.I.G.
- Colquitt, J., Royle, P. & Waugh, N. Are analogue insulins better than soluble in continuous subcutaneous insulin infusion? Results of a meta-analysis, 863
- Conget, I. *see* Costa, A.
- Connolly, V.M. *see* Vaidya, B.
- Connor, H. *see* Dang, C.N.
- Coppini, D.V. *see* Spruce, M.C.
- Costa, A., Casamirjana, R., Casals, E., Álvarez, L., Morales, J., Masramón, X., Hernández, G., Gomis, R. & Conget, I. Effects of atorvastatin on glucose homeostasis, postprandial triglyceride response and C-reactive protein in subjects with impaired fasting glucose, 743
- de Courten, M.P. *see* Tapp, R.J.
- de Courten, M.P. *see* Williams, J.W.
- Cox, J. *see* Valabhji, J.
- Craft, A.W. *see* Parker, L.
- Critchley, J.A.J.H. *see* Thomas, G.N.
- Croall, J. *see* Hassanein, M.M.
- Croft, D.J. *see* Hadden, D.R.
- Cromme, P. *see* Schramm, W.
- Crosland, A. *see* Whitford, D.L.
- Cross, D. *see* Zhao, H.X.
- Cruickshank, J.K. *see* Vyas, A.
- Cull, C.A. *see* Davis, T.M.E.
- Cull, C.A. *see* Hadden, D.R.
- Cundy, T. & Drury, P. Insulin-treated diabetes and driving in the UK, 168
- Czupryniak, L., Ruxer, J., Saryusz-Wolska, M. & Loba, J. Discontinuing insulin therapy after diabetic ketoacidosis—is it cause worth considering? 333
- Dalfollo, M. *see* Inchiostro, S.
- Damm, P. *see* Jensen, D.M.
- Dang, C. *see* Armstrong, D.G.
- Dang, C.N., Prasad, Y.D.M., Boulton, A.J.M. & Jude, E.B. Methicillin-resistant *Staphylococcus aureus* in the diabetic foot clinic: a worsening problem, 159
- Dang, C.N., Prasad, Y.D.M., Connor, H. & Boulton, A.J.M. Update on the Diabetic Foot from Malvern Foot Conference 2002, 245
- Davey Smith, G. *see* Bachmann, M.O.
- Davies, H.A. *see* Patel, V.K.
- Davies, S. *see* Rauce, J.
- Davis, E.A. *see* Caplin, N.J.
- Davis, T.M.E., Mehta, Z., Mackay, I.R., Cull, C.A., Bruce, D.G., Fida, S., Rowley, M.J. & Holman, R.R. Autoantibodies to the islet cell antigen SOX-13 are associated with duration but not type of diabetes, 198
- De Venuto, G. *see* Inchiostro, S.
- Dean, H.J. *see* Kaila, B.
- Deary, I.J. *see* Henderson, J.N.
- Deary, I.J. *see* McCrimmon, R.J.
- Deary, I.J. *see* Sommerfield, A.J.
- Deary, I.J. *see* Strachan, M.W.J.
- Deary, I.J. *see* Taylor, M.D.
- Deepa, R. *see* Velmurugan, K.
- Dekker, J.M. *see* Adriaanse, M.C.
- Dekker, J.M. *see* Scheffer, P.G.
- Delaney, P. *see* Trembl, H.
- Delgado, E. *see* Botas, P.
- Demaine, A.G. *see* Zhao, H.X.
- De Valk, H.W. *see* Goudswaard, A.N.
- DeVries, J.H., Lindholm, A., Jacobsen, J.L., Heine, R.J. & Home, P.D. The Tri-Continental Insulin Aspart Study Group. A randomized trial of insulin aspart with intensified basal NPH insulin supplementation in people with Type 1 diabetes, 312
- Dhindsa, P., Scott, A.R. & Donnelly, R. Metabolic and cardiovascular effects of very-low-calorie diet therapy in obese patients with Type 2 diabetes in secondary failure: outcomes after 1 year, 319
- Díaz-Cadorniga, F.J. *see* Botas, P.
- Díaz de Greñu, C. *see* Botas, P.
- Dickinson, H.O. *see* Parker, L.
- Diebler, F. *see* Roach, P.
- Dinneen, S.F. *see* Yuyun, M.F.
- Dixon, A.F. *see* Dixon, J.B.
- Dixon, J.B., Dixon, A.F., & O'Brien, P.E. Improvements in insulin sensitivity and  $\beta$ -cell function (HOMA) with weight loss in the severely obese, 127
- Doll, H. *see* Farmer, A.J.
- Dong, L. *see* Yang, X.
- Donnelly, R. *see* Dhindsa, P.
- Donohoe, M. *see* Owen, K.R.
- Dornan, T. *see* Carroll, C.
- Dornan, T. & Carroll, C. Medical communication and diabetes, 85
- Dornan, T.L. *see* Kaushal, K.
- Dreyer, M. *see* Roach, P.
- Drury, P. *see* Cundy, T.
- Drury, P.L. *see* Manning, P.J.
- Dudenhausen, J.W. *see* Schaefer-Graf, U.M.
- Duffy, S.W. *see* Liu, W.-J.

- Dunger, D.B. *see* Ford-Adams, M.E.
- Dunne, F., Brydon, P., Smith, K. & Gee, H. Pregnancy in women with Type 2 diabetes: 12 years outcome data 1990–2002, 734
- Dunne, C.J. *see* Mazze, R.S.
- Dunne, F. *see* Sinha, B.
- Dunne, F.P. *see* Chatterjee, S.
- Dunstan, D.W. *see* Tapp, R.J.
- Dyer, P.H. *see* Bain, S.C.
- Eachus, J. *see* Bachmann, M.O.
- Ebrahim, S. *see* Bott, U.
- Edenborough, F.P. *see* Mackie, A.D.R.
- Edgar, K. *see* Skinner, T.C.
- Edge, J.A. *see* Ford-Adams, M.E.
- Edwards, O.M. *see* Yuyun, M.F.
- Egberts, E.-H. *see* Holstein, A.
- Eguchi, H. *see* Marugame, T.
- Eichbauer-Sturm, G. *see* Biesenbach, G.
- Eiger, G. *see* Kian, K.
- van Eijk, J.Th.M. *see* Renders, C.M.
- ElBouri, K. *see* Prakash, P.K.
- Elkeles, R.S. *see* Valabhji, J.
- Ellard, S. *see* Owen, K.R.
- Elwig, C. *see* Valabhji, J.
- Enas, E.A. *see* Velmurugan, K.
- Erginay, A. *see* Massin, P.
- Eriksson, J.W. *see* Lindmark, S.
- Ernst, E. *see* Schmidt, K.
- Espiner, E.A. *see* Manning, P.J.
- Evans, J. *see* Tang, E.W.
- Ewing, F.M.E. *see* McCrimmon, R.J.
- Ewing, F.M.E. *see* Sommerfield, A.J.
- Ewins, D.L. *see* Hassanein, M.M.
- Fait, D. *see* Inchiostro, S.
- Fall, C.H.D. *see* Newsome, C.A.
- Farmer, A.J., Doll, H., Levy, J.C. & Salkovskis, P.M. The impact of screening for Type 2 diabetes in siblings of patients with established diabetes, 996
- Feest, T. *see* Schramm, W.
- Feher, M.D. *see* Trembl, H.
- Feldt-Rasmussen, B. *see* Schramm, W.
- Feltbower, R.G. *see* McKinney, P.A.
- Feltbower, R.G., McKinney, P.A., Parslow, R.C., Stephenson, C.R. & Bodansky, H.J. Type 1 diabetes in Yorkshire, UK: time trends in 0–14 and 15–29-year-olds, age at onset and age-period-cohort modelling, 437
- Feskens, E.J.M. *see* Mensink, M.
- Fida, S. *see* Davis, T.M.E.
- Fiestras-Janeiro, G. *see* García-Estévez, D.A.
- Firth, R.G. *see* Smith, S.M.
- Fischbacher, C.M., Bhopal, R., Rutter, M.K., Unwin, N.C., Marshall, S.M., White, M. & Alberti, K.G.M.M. Microalbuminuria is more frequent in South Asian than in European origin populations: a comparative study in Newcastle, UK, 31
- Fisher, M. *see* Winocour, P.H.
- Florkowski, C.M., Scott, R.S., Graham, P.J., Han, D.Y. & Moir, C.L. Cause-specific and total mortality in the Canterbury (New Zealand) insulin-treated Diabetic Registry population: a 15-year follow-up study, 191
- Flyvbjerg, A. *see* Frystyk, J.
- Ford-Adams, M.E., Murphy, N.P., Moore, E.J., Edge, J.A., Ong, K.L., Watts, A.P., Acerini, C.L. & Dunger, D.B. Insulin lispro: a potential role in preventing nocturnal hypoglycaemia in young children with diabetes mellitus, 656
- Forrest, K.Y.-Z. *see* Bosnyak, Z.
- Forrester, J.V. *see* Olson, J.A.
- Foy, C. *see* Scanlon, P.H.
- Frankel, S. *see* Bachmann, M.O.
- Frier, B.M. *see* Henderson, J.N.
- Frier, B.M. *see* McCrimmon, R.J.
- Frier, B.M. *see* Sommerfield, A.J.
- Frier, B.M. *see* Strachan, M.W.J.
- Frier, B.M. *see* Taylor, M.D.
- Frykberg, R.G. *see* Armstrong, D.G.
- Frystyk, J., Bek, T., Flyvbjerg, A., Skjærbaek, C. & Ørskov, H. The relationship between the circulating IGF system and the presence of retinopathy in Type 1 diabetic patients, 269
- Furler, S.M. *see* Kriketos, A.D.
- Furlong, N.J., Hulme, S.A., O'Brien, S.V. & Hardy, K.J. Comparison of repaglinide vs. gliclazide in combination with bedtime NPH insulin in patients with Type 2 diabetes inadequately controlled with oral hypoglycaemic agents, 935
- Gallen, I.W. *see* Masson, E.A.
- Gallois, Y. *see* Hadjadj, S.
- Ganss, R. *see* Birkeland, K.I.
- García-Estévez, D.A., Araújo-Vilar, D., Fiestras-Janeiro, G., Saavedra-Gonzalez, Á. & Cabezas-Cerrato, J. Insulin resistance in essential hypertension: a conflictive point of view, 1035
- Gaudric, A. *see* Massin, P.
- Gee, H. *see* Chatterjee, S.
- Gee, H. *see* Dunne, F.
- Gentile, L. *see* Bo, S.
- Gibson, J. *see* Harding, S.
- Gibson, J.M. *see* Kaushal, K.
- Gilbert, R.E. *see* Krum, H.
- Gilbert, R.E., Krum, H., Wilkinson-Berka, J. & Kelly, D.J. The renin-angiotensin system and the long-term complications of diabetes: pathophysiological and therapeutic considerations, 607
- Gill, G.V. *see* Bain, S.C.
- Gill, S. *see* Vyas, A.
- Giorgino, F. *see* Greco, P.
- Glümer, C. *see* Borch-Johnsen, K.
- Glümer, C. *see* Colagiuri, S.
- Goatman, K.A. *see* Olson, J.A.
- Goddard, J.R. *see* Holt, R.I.G.
- Gold, A.E. *see* McCrimmon, R.J.
- Gold, A.E. *see* Taylor, M.D.
- Gomis, R. *see* Costa, A.
- Gone, K. *see* Barton, D.M.
- Gottsäter, A., Rydén-Ahlgren, Å., Szegla, B., Hedblad, B., Persson, J., Berglund, G., Wroblewski, M. & Sundkvist, G. Cardiovascular autonomic neuropathy associated with carotid atherosclerosis in Type 2 diabetic patients, 495
- Goudswaard, A.N., Stolk, R.P., de Valk, H.W. & Rutten, G.E.H.M. Improving glycaemic control in patients with Type 2 diabetes mellitus without insulin therapy, 540
- Gough, K. *see* Strachan, M.W.J.
- Gouws, E. *see* Motala, A.A.

- Graham, P.J. *see* Florkowski, C.M.
- Grant, P.J. *see* Barnett, A.H.
- Grant, P.J. *see* Kain, K.
- Gray, A. *see* Clarke, P.
- Greco, P., Vimercati, A., Hyett, J., Rossi, A.C., Scioscia, M., Giorgino, F., Loverro, G. & Selvaggi, L. The ultrasound assessment of adipose tissue deposition in fetuses of 'well controlled' insulin-dependent diabetic pregnancies, 858
- Greene, S. *see* Skinner, T.C.
- Greenwood, R. *see* Harding, S.
- Gregory, J.W. *see* Channon, S.
- Griffin, S.J. *see* Rahman, M.
- Gudat, U. *see* Roach, P.
- Guillausseau, P.J. *see* Massin, P.
- Hadden, D.R., Cull, C.A., Croft, D.J. & Holman, R.R. Poor pregnancy outcome for women with Type 2 diabetes, 506
- Hadjadj, S., Gallois, Y., Alhenc-Gelas, F., Chatellier, G. & Marre, M. for the DZABHYCAR Study Group. Angiotensin-I-converting enzyme insertion/deletion polymorphism and high urinary albumin concentration in French Type 2 diabetes patients, 677
- Hahn, J.S. *see* Kim, D.J.
- Haideri, A.Z. *see* Vyas, A.
- Hampson, S.E. *see* Clark, M.
- Hamulyak, K. *see* Leurs, P.B.
- Han, D.Y. *see* Florkowski, C.M.
- Hanna, F.W. *see* Prakash, P.K.
- Hanrath, P. *see* Ortlepp, J.R.
- Hansen, K.W., Poulsen, P.L. & Mogensen, C.E. Ambulatory blood pressure monitoring, 509
- Hansen, S.K. *see* Pildal, J.
- Hansen, T. *see* Pildal, J.
- Hanssen, K.F. *see* Birkeland, K.I.
- Harding, S., Greenwood, R., Aldington, S., Gibson, J., Owens, D., Taylor, R., Kohner, E., Scanlon, P. & Leese, G. The Diabetic Retinopathy Grading and Disease Management Working Party. Grading and disease management in national screening for diabetic retinopathy in England and Wales, 965
- Harding, S.P. *see* Younis, N.
- Hardy, K.J. *see* Furlong, N.J.
- Hardy, K.J. *see* Moulik, P.K.
- Harney, B. *see* Scanlon, P.H.
- Harvey, R. *see* Morris, A.
- Hassanein, M.M., Croall, J., Ewins, D.L. & Worth, R.C. An unusual cause of diabetic ketoacidosis and fulminant septicaemia, 242
- Hattersley, A.T. *see* Owen, K.R.
- Hattori, N., Rühl, J., Bengel, F.M., Nekolla, S.G., Standl, E., Schwaiger, M. & Schnell, O. Cardiac autonomic dysinnervation and myocardial blood flow in long-term Type 1 diabetic patients, 375
- Hayasaka, H. *see* Marugame, T.
- Hedblad, B. *see* Gottsäter, A.
- Heine, R.J. *see* Adriaanse, M.C.
- Heine, R.J. *see* DeVries, J.H.
- Heine, R.J. *see* Renders, C.M.
- Heine, R.J. *see* Scheffer, P.G.
- Heinze, T. *see* Schaefer-Graf, U.M.
- Heise, T. on behalf of the authors. Reply, 334
- Heller, S. The National Service Framework for Diabetes: a glass half empty or half full? 173
- Henderson, J.N., Allen, K.V., Deary, I.J. & Frier, B.M. Hypoglycaemia in insulin-treated Type 2 diabetes: frequency, symptoms and impaired awareness, 1016
- Henrich, W. *see* Schaefer-Graf, U.M.
- Hepburn, D.A. *see* McCrimmon, R.J.
- Hermann, N. *see* Nordly, S.
- Hermus, A.R. *see* Pouwels, M.-J.J.
- Hernández, G. *see* Costa, A.
- Hesketh, K. *see* Cameron, F.J.
- Hey, S. *see* Viner, R.M.
- Hipwell, J.H. *see* Olson, J.A.
- Hirasawa, K. *see* Yokoyama, H.
- Hirschberger, S. *see* Bott, U.
- Hitman, G.A. *see* Barnett, A.H.
- Ho, L.-T. *see* Pan, W.-H.
- Hod, M. *see* Yorgev, Y.
- Hödl, R. *see* Siebenhofer, A.
- HOE 901/2004 Study Investigators Group. Safety and efficacy of insulin glargine (HOE 901) versus NPH insulin in combination with oral treatment in Type 2 diabetic patients, 545
- Hoffmann, R. *see* Ortlepp, J.R.
- Hojlund, K. *see* Staehr, P.
- Holdaway, I.M. *see* Manning, P.J.
- Hollins, A. *see* Sinha, B.
- Holman, R. *see* Clarke, P.
- Holman, R.R. *see* Davis, T.M.E.
- Holman, R.R. *see* Hadden, D.R.
- Holohan, J. *see* Smith, S.M.
- Holst, J.J. *see* Staehr, P.
- Holstein, A., Plaschke, A. & Egberts, E.-H. Use of insulin glargine during embryogenesis in a pregnant woman with Type 1 diabetes, 779
- Holt, R.I.G., Goddard, J.R., Clarke, P. & Coleman, M.A.G. A postnatal fasting plasma glucose is useful in determining which women with gestational diabetes should undergo a postnatal oral glucose tolerance test, 594
- Holt, R.I.G., Simpson, H.L. & Sönksen, P.H. The role of the growth hormone-insulin-like growth factor axis in glucose homeostasis, 3
- Home, P.D. *see* DeVries, J.H.
- Hopkins, D. *see* Bingham, E.
- Hopper, C.D. *see* Bachmann, M.O.
- Hother-Nielsen, O. *see* Staehr, P.
- Hounsborne, B. *see* Kaushal, K.
- Howard, E.R. *see* Watkins, P.J.
- Howells, L. *see* Skinner, T.C.
- Hsu-Hage, B.H.-H. *see* Yang, X.
- Hughes, P. *see* Zhao, H.X.
- Huh, K.B. *see* Kim, D.J.
- Hulme, S.A. *see* Furlong, N.J.
- Hwu, C.-M. *see* Pan, W.-H.
- Hyett, J. *see* Greco, P.
- Idrogo, M.A. *see* Mazze, R.S.
- Illahi, W. *see* Basu, A.
- Inagaki, Y. *see* Pang, H.
- Inchiostro, S., Dalfollo, M., Marzano, A., Citroni, N., Peccatori, S., Fait, D. & De Venuto, G. Prevalence of diabetes and/or ischaemic heart disease in classes of increasing carotid artery atherosclerosis: an ultrasonographic study, 670



- Ingberg, C.-M., Särnblad, S., Palmér, M., Schvarcz, E., Berne, C. & Åman, J. Body composition in adolescent girls with Type 1 diabetes, 1005
- Inukai, T. *see* Aso, Y.
- Ishiyama, M. *see* Yokoyama, H.
- Jacobsen, J.L. *see* DeVries, J.H.
- Jácome de Castro, J. *see* Macedo, A.
- Jagannathan, N.R. *see* Misra, A.
- Jardine, D.L. *see* Tang, E.W.
- Jenkins, A.B. *see* Kriketos, A.D.
- Jenkins, D. *see* Sinha, B.
- Jensen, D.M., Damm, P., Sørensen, B., Mølsted-Pedersen, L., Westergaard, J.G., Korsholm, L., Ovesen, P. & Beck-Nielsen, H. Proposed diagnostic thresholds for gestational diabetes mellitus according to a 75-g oral glucose tolerance test. Maternal and perinatal outcomes in 3260 Danish women, 51
- Jick, H. *see* Wood, S.L.
- Johansson, A. *see* Skinner, T.C.
- Jones, A.F. *see* Al-Daghri, N.
- Jones, A.F. *see* Bain, S.C.
- Jones, K.E. *see* Bain, S.C.
- Jones, T.W. *see* Caplin, N.J.
- Jorge, Z. *see* Macedo, A.
- Jørgensen, T. *see* Nordly, S.
- Jude, E.B. *see* Dang, C.N.
- Kaila, B., Dean, H.J., Schroeder, M. & Taback, S.P. HLA, day care attendance, and socio-economic status in young patients with Type 1 diabetes, 777
- Kain, K., Catto, A.J. & Grant, P.J. Associations between insulin resistance and thrombotic risk factors in high-risk South Asian subjects, 651
- Kamal, A.D. *see* Basu, A.
- Kamalakkannan, D. *see* Barton, D.M.
- Kanda, T. *see* Aso, Y.
- Kaplan, B. *see* Yoge, Y.
- Kaushal, K. & Dornan, T.L. Evidence-Based Diabetes Care, 510
- Kaushal, K., Gibson, J.M., Railton, A., Hounscome, B., New, J.P. & Young, R.J. A protocol for improved glycaemic control following corticosteroid therapy in diabetic pregnancies, 73
- Kearney, M.T. *see* Wheatcroft, S.B.
- Kelly, D.J. *see* Gilbert, R.E.
- Kelly, W.F. *see* Vaidya, B.
- Kendall, D. *see* Lunt, H.
- Kendall, D.M. *see* Mazze, R.S.
- Khan, M. *see* Basu, A.
- Kian, K. & Eiger, G. Anticoagulant therapy in hyperosmolar non-ketotic diabetic coma, 603
- Kilhovd, B. *see* Birkeland, K.I.
- Kim, D.J., Kim, Y.M., Yun, Y.S., Ahn, C.W., Cha, B.S., Song, Y.D., Lim, S.K., Kim, K.R., Hahn, J.S., Huh, K.B. & Lee, H.C. Therapeutic effect of recombinant human erythropoietin on anaemia with erythropoietin deficiency in diabetic patients, 661
- Kim, D.R. *see* Choi, K.M.
- Kim, K.B. *see* Choi, K.M.
- Kim, K.R. *see* Kim, D.J.
- Kim, N.H. *see* Choi, K.M.
- Kim, S.K. *see* Choi, K.M.
- Kim, Y.M. *see* Kim, D.J.
- Kimura, H. *see* Pang, H.
- Kirkpatrick, J.N. *see* Scanlon, P.H.
- Kjos, S.L. *see* Schaefer-Graf, U.M.
- Klepper, J. *see* Willemsen, M.A.A.P.
- Ko, G.T.C., Lee, S.C., Pu, Y.B., Ng, M.C.Y., So, W.-Y., Thomas, N., Chan, W.B., Cockram, C.S. & Chan, J.C.N. Tumour necrosis factor-alpha promoter gene polymorphism at -308 (genotype AA) in Chinese subjects with Type 2 diabetes, 167
- Kobayashi, I. *see* Aso, Y.
- Koda, Y. *see* Pang, H.
- Kohner, E. *see* Harding, S.
- Konopatskaya, O., Whatmore, J.L., Tooke, J.E. & Shore, A.C. Insulin and lysophosphatidylcholine synergistically stimulate NO-dependent cGMP production in human endothelial cells, 838
- von Korff, A. *see* Ortlepp, J.R.
- Korsholm, L. *see* Jensen, D.M.
- Koyama, K. *see* Yokoyama, H.
- Kriegsman, D.M.W. *see* Renders, C.M.
- Kriketos, A.D., Carey, D.G., Jenkins, A.B., Chisholm, D.J., Furler, S.M. & Campbell, L.V. Central fat predicts deterioration of insulin secretion index and fasting glycaemia: 6-year follow-up of subjects at varying risk of Type 2 diabetes mellitus, 294
- Kruishoop, M. *see* Mensink, M.
- Krum, H. *see* Gilbert, R.E.
- Krum, H., Skiba, M. & Gilbert, R.E. Comparative metabolic effects of hydrochlorothiazide and indapamide in hypertensive diabetic patients receiving ACE inhibitor therapy, 708
- Krzentowski, G. *see* Zhang, L.
- Kumar, M. *see* Misra, A.
- Kumar, S. *see* Al-Daghri, N.
- Lacerda Nobre, E. *see* Macedo, A.
- Lahousen, T., Lipp, R.W. & Schnedl, W.J. Can hyperglycaemia cause jejuno-jejunal intussusception?, 335
- Lajer, M. *see* Pildal, J.
- Lamont, D.W. *see* Parker, L.
- Lamont, S.S. *see* Whitford, D.L.
- Landgraf, R. *see* Schramm, W.
- Lauritzen, T. *see* Borch-Johnsen, K.
- Lauritzen, T. *see* Colagiuri, S.
- Law, C.M. *see* Newsome, C.A.
- Lawrence, J.B. *see* Velmurugan, K.
- Lean, M.E.J. The implementation of nutritional advice for people with diabetes, 783
- Lee, H.C. *see* Kim, D.J.
- Lee, J. *see* Choi, K.M.
- Lee, K. *see* Marugame, T.
- Lee, L.-T. *see* Liu, W.-J.
- Lee, S.C. *see* Ko, G.T.C.
- Leese, G. *see* Harding, S.
- Leese, G. *see* Morris, A.
- Leese, G.P., Morris, A.D. & Olson, J. A national retinal screening programme for diabetes in Scotland, 962
- Lefèbvre, P.J. *see* Zhang, L.
- Legood, R. *see* Clarke, P.
- Leurs, P.B., van Oerle, R., Hamulyak, K. & Wolffenbuttel, B.H.R. Tissue factor pathway inhibitor (TFPI) release after heparin stimulation is increased in Type 1 diabetic patients with albuminuria, 16
- Levy, J.C. *see* Farmer, A.J.

- Lewis-Barned, N. *see* Scanlon, P.H.
- Lezo, A. *see* Bo, S.
- Lim, S.K. *see* Kim, D.J.
- Lindholm, A. *see* DeVries, J.H.
- Lindmark, S., Wiklund, U., Björle, P. & Eriksson, J.W. Does the autonomic nervous system play a role in the development of insulin resistance? A study on heart rate variability in first-degree relatives of Type 2 diabetes patients and control subjects, 399
- Lindow, S.W. *see* Masson, E.A.
- Lindström, J. *see* Qiao, Q.
- Lins, P.-E. *see* Bragd, J.
- Lipkin, G. *see* Chatterjee, S.
- Lipp, R.W. *see* Lahousen, T.
- Liu, W.-J., Lee, L.-T., Yen, M.-F., Tung, T.-H., Williams, R., Duffy, S.W. & Chen, T.H.-H. Assessing progression and efficacy of treatment for diabetic retinopathy following the proliferative pathway to blindness: implication for diabetic retinopathy screening in Taiwan, 727
- Loba, J. *see* Czupryniak, L.
- Lovelock, L. *see* Arun, C.S.
- Loverro, G. *see* Greco, P.
- Lunt, H., Kendall, D., Anderson, P., Willis, J.A. & Moore, M.P. Prevalence of microvascular complications in adolescents with Type 1 diabetes, 421
- Lutterman, J.A. *see* Pouwels, M.-J.J.
- Łysiak-Szydłowska, W. *see* Rachoń, D.
- Macdonald, I.A. *see* Bingham, E.
- Macedo, A., Jorge, Z., Lacerda Nobre, E., Pratas, S. & Jácome de Castro, J. Prevalence of Type 1 diabetes mellitus in Portugal, 1995–1999: cohort of young men, 418
- Mackay, I.R. *see* Davis, T.M.E.
- Mackie, A.D.R., Thornton, S.J. & Edenborough, F.P. Cystic fibrosis-related diabetes, 425
- MacLeod, K.M. *see* McCrimmon, R.J.
- Mair, G. *see* Penney, G.C.
- Malhotra, R. *see* Scanlon, P.H.
- Manfrini, S. *see* Pozzilli, P.
- Mankovsky, B.N., Piolot, R., Mankovsky, O.L. & Ziegler, D. Impairment of cerebral autoregulation in diabetic patients with cardiovascular autonomic neuropathy and orthostatic hypotension, 119
- Mankovsky, O.L. *see* Mankovsky, B.N.
- Manning, P.J., Espiner, E.A., Yoon, K., Drury, P.L., Holdaway, I.M. & Bowers, A. An unusual cause of hyperinsulinaemic hypoglycaemia syndrome, 772
- Mannucci, E., Bardini, G., Rotella, F. & Rotella, C.M. Comparison among different insulin sensitivity indices in obese patients, 462
- Marchisio, B. *see* Bo, S.
- Marre, M. *see* Hadjadj, S.
- Marre, M. *see* Massin, P.
- Marsden, P. *see* Carroll, C.
- Marshall, S.M. *see* Fischbacher, C.M.
- Marugame, T., Hayasaka, H., Lee, K., Eguchi, H. & Matsumoto, S. Alveolar bone loss associated with glucose tolerance in Japanese men, 746
- Marzano, A. *see* Inchiostro, S.
- Maser, R.E. *see* Bosnyak, Z.
- Mason, J. National screening for diabetic retinopathy: clear vision needed, 959
- Masramón, X. *see* Costa, A.
- Massin, P., Erginay, A., Ben Mehidi, A., Vicaute, E., Quentel, G., Victor, Z., Marre, M., Guillausseau, P.J. & Gaudric, A. Evaluation of a new non-mydiatic digital camera for detection of diabetic retinopathy, 635
- Massobrio, M. *see* Bo, S.
- Masson, E.A., Patmore, J.E., Brash, P.D., Baxter, M., Caldwell, G., Gallen, I.W., Price, P.A., Vice, P.A., Walker, J.D. & Lindow, S.W. Pregnancy outcome in Type 1 diabetes mellitus treated with insulin lispro (Humalog), 46
- Mather, H. *see* Barnett, A.H.
- Mathers, J.C. *see* Parker, L.
- Matsumoto, S. *see* Marugame, T.
- Mattock, M.B. *see* Pickup, J.C.
- Mazze, R.S., Simonson, G.D., Robinson, R.L., Kendall, D.M., Idrogo, M.A., Adlis, S.A., Boyce, K.S., Dunne, C.J., Anderson, R.L. & Bergenstal, R.M. Characterizing blood pressure control in individuals with Type 2 diabetes: the relationship between clinic and self-monitored blood pressure, 752
- McAuliffe, A. *see* Smith, S.M.
- McCrimmon, R.J., Deary, I.J., Gold, A.E., Hepburn, D.A., MacLeod, K.M., Ewing, F.M.E. & Frier, B.M. Symptoms reported during experimental hypoglycaemia: effect of method of induction of hypoglycaemia and of diabetes per se, 507
- McCulloch, A.J. *see* Razvi, S.
- McDonald, D. *see* Moore, K.B.
- McEvelly, A. *see* Skinner, T.C.
- McFarlane, S.I., Shin, J., Byrne, K. & Williams, R. Reply, 336
- McHardy, K.C. *see* Olson, J.A.
- McKenna, K. *see* Moore, K.B.
- McKibbin, M.A. *see* Agrawal, A.
- McKinney, P.A., Feltbower, R.G. & Bodansky, H.J. Reply, 602
- McKinney, P.A. *see* Feltbower, R.G.
- McKnight, J.A. *see* Strachan, M.W.J.
- Mehta, Z. *see* Davis, T.M.E.
- Meiselman, H.J. *see* Chong-Martinez, B.
- Menato, G. *see* Bo, S.
- Mensink, M., Feskens, E.J.M., Kruijschoop, M., de Bruin, T.W.A., Saris, W.H.M. & Blaak, E.E. Subscapular skinfold thickness distinguishes between transient and persistent impaired glucose tolerance: Study on Lifestyle-Intervention and Impaired Glucose Tolerance Maastricht (SLIM), 552
- Metrikat, J. *see* Ortlepp, J.R.
- Millward, B.A. *see* Zhao, H.X.
- Misra, A., Sinha, S., Kumar, M., Jagannathan, N.R. & Pandey, R.M. Proton magnetic resonance spectroscopy study of soleus muscle in non-obese healthy and Type 2 diabetic Asian Northern Indian males: high intramyocellular lipid content correlates with excess body fat and abdominal obesity, 361
- Mogensen, C.E. *see* Hansen, K.W.
- Mohan, V. *see* Velmurugan, K.
- Moir, C.L. *see* Florkowski, C.M.
- Mølsted-Pedersen, L. *see* Jensen, D.M.
- Moore, E.J. *see* Ford-Adams, M.E.
- Moore, K.B., McKenna, K., Tormey, W.P., McDonald, D. & Thompson, C.J. The albuminuric action of atrial natriuretic peptide is not modified by ACE-inhibition with perindopril in Type 2 diabetes, 713
- Moore, M.P. *see* Lunt, H.
- Morales, J. *see* Costa, A.

- Morris, A., Slattery, J., Olson, J., Leese, G. & Harvey, R.  
Current status of screening for diabetic retinopathy in the UK, 603
- Morris, A.D. *see* Leese, G.P.
- Mortensen, H.B. *see* Nordly, S.
- Motala, A.A., Pirie, F.J., Gouws, E., Amod, A. & Omar, M.A.K.  
High incidence of Type 2 diabetes mellitus in South African Indians: a 10-year follow-up study, 23
- Moulik, P.K. & Hardy, K.J. Hypertension, anti-hypertensive drug therapy and erectile dysfunction in diabetes, 290
- Mukherjee, S. *see* Sharp, P.S.
- Munro, A. *see* Sinha, B.
- Murphy, M. *see* Bain, S.C.
- Murphy, N.P. *see* Ford-Adams, M.E.
- Myers, L. *see* Razvi, S.
- Myśliwska, J. *see* Rachoń, D.
- Nagai, Y., Nakatsumi, Y., Abe, T. & Nomura, G. Is the severity of obstructive sleep apnoea associated with the degree of insulin resistance? 81
- Nagai, Y., Sugiyama, Y., Nakatsumi, Y., Abe, T. & Nomura, G.  
Rapid amelioration of muscle sympathetic nerve activity by pioglitazone in an obese Type 2 diabetic patient, 868
- Nakatsumi, Y. *see* Nagai, Y.
- Naylor, E. *see* Carroll, C.
- Neal, R.D. *see* Alazri, M.H.
- Nekkolla, S.G. *see* Hattori, N.
- New, J.P. *see* Kaushal, K.
- New, J.P. *see* Tieszen, K.L.
- Newsome, C.A., Shiell, A.W., Fall, C.H.D., Phillips, D.I.W., Shier, R. & Law, C.M. Is birth weight related to later glucose and insulin metabolism?—a systematic review, 339
- Ng, L.L. *see* Siebenhofer, A.
- Ng, M.C.Y. *see* Ko, G.T.C.
- Ngugi, N. *see* Arun, C.S.
- Nijpels, G. *see* Adriaanse, M.C.
- Nixon, B.P. *see* Armstrong, D.G.
- Nixon, S.J. *see* Saeed, B.O.
- Nomura, G. *see* Nagai, Y.
- Nordly, S., Jørgensen, T., Andreasen, A.H., Hermann, N. & Mortensen, H.B. Quality of diabetes management in children and adolescents in Denmark, 568
- Nørgaard, K. A nationwide study of continuous subcutaneous insulin infusion (CSII) in Denmark, 307
- Nomura, G. *see* Nagai, Y.
- Northam, E.A. *see* Cameron, F.J.
- Nutrition Subcommittee of the Diabetes Care Advisory Committee of Diabetes UK. The implementation of nutritional advice for people with diabetes, 786
- O'Brien, P.E. *see* Dixon, J.B.
- O'Brien, S.V. *see* Furlong, N.J.
- O'Leary, M. *see* Smith, S.M.
- O'Leary, P. *see* Caplin, N.J.
- O'Malley, B. *see* Rance, J.
- van Oerle, R. *see* Leurs, P.B.
- Okamoto, T. *see* Pang, H.
- Okumura, K. *see* Aso, Y.
- Olson, J. *see* Leese, G.P.
- Olson, J. *see* Morris, A.
- Olson, J.A., Strachan, F.M., Hipwell, J.H., Goatman, K.A., McHardy, K.C., Forrester, J.V. & Sharp, P.F. A comparative evaluation of digital imaging, retinal photography and optometrist examination in screening for diabetic retinopathy, 528
- Omar, M.A.K. *see* Motala, A.A.
- Ong, K.L. *see* Ford-Adams, M.E.
- Orchard, T.J. *see* Bosnyak, Z.
- Ørskov, H. *see* Frystyk, J.
- Ortlepp, J.R., Metrikat, J., Albrecht, M., von Korff, A., Hanrath, P. & Hoffmann, R. The vitamin D receptor gene variant and physical activity predicts fasting glucose levels in healthy young men, 451
- Osawa, K. *see* Ota, T.
- Oskarsson, P. *see* Bragð, J.
- Ota, T. & Osawa, K. Marked improvement of glycaemic control with pioglitazone in a Type 2 diabetic patient associated with Charcot-Marie-Tooth disease, 420
- Ovesen, P. *see* Jensen, D.M.
- Owen, K.R., Donohoe, M., Ellard, S. & Hattersley, A.T. Response to treatment with rosiglitazone in familial partial lipodystrophy due to a mutation in the LMNA gene, 823
- Owen, K.R., Roland, J., Smith, K. & Hattersley, A.T. Adolescent onset Type 2 diabetes in a non-obese Caucasian patient with an unbalanced translocation, 483
- Owens, D. *see* Harding, S.
- Owens, D.R., Zinman, B. & Bolli, G. Alternative routes of insulin delivery, 886
- Padfield, P.L. *see* Strachan, M.W.J.
- Pagano, G. *see* Bo, S.
- Palmér, M. *see* Ingberg, C-M.
- Pan, W.-H., Yeh, W.-T., Chang, H.-Y., Hwu, C.-M. & Ho, L.-T.  
Prevalence and awareness of diabetes and mean fasting glucose by age, sex, and region: results from the Nutrition and Health Survey in Taiwan, 1993–1996, 182
- Pandey, R.M. *see* Misra, A.
- Pang, H., Koda, Y., Yamagishi, S.-I., Amano, S., Inagaki, Y., Okamoto, T., Yamada, K. & Kimura, H. N-acetylglucosamine-phosphate mutase genotype and diabetic microvascular complications, 419
- Park, I.B. *see* Choi, K.M.
- Parker, L., Lamont, D.W., Unwin, N., Pearce, M.S., Bennett, S.M.A., Dickinson, H.O., White, M., Mathers, J.C., Alberti, K.G.M.M. & Craft, A.W. A lifecourse study of risk for hyperinsulinaemia, dyslipidaemia and obesity (the central metabolic syndrome) at age 49–51 years, 406
- Parkin, T. & Skinner, T.C. Discrepancies between patient and professionals recall and perception of an outpatient consultation, 909
- Parslow, R.C. *see* Feltbower, R.G.
- Patel, V.K. & Davies, H.A. Insulin resistance Type A and short 5th metacarpals, 500
- Patmore, J.E. *see* Masson, E.A.
- Patton, K. *see* Razvi, S.
- Pawa, M. *see* Barnett, A.H.
- Pearce, M.S. *see* Parker, L.
- Pearson, D.W.M. *see* Penney, G.C.
- Pearson, N.J. *see* Bachmann, M.O.
- Peccatori, S. *see* Inchiostro, S.



- Pedersen, O. *see* Pildal, J.  
 Pein, M. *see* Roach, P.  
 Penney, G.C., Mair, G. & Pearson, D.W.M. The relationship between birth weight and maternal glycated haemoglobin (HbA<sub>1c</sub>) concentration in pregnancies complicated by Type 1 diabetes, 162  
 Pernet, A. *see* Bingham, E.  
 Persson, J. *see* Gottsäter, A.  
 Pfohl, U. *see* Zimny, S.  
 Phillip, M. *see* Yogeve, Y.  
 Phillips, C. *see* Rance, J.  
 Phillips, D.I.W. *see* Newsome, C.A.  
 Picardi, A. *see* Pozzilli, P.  
 Pickup, J.C. & Mattock, M.B. Activation of the innate immune system as a predictor of cardiovascular mortality in Type 2 diabetes mellitus, 723  
 Pickup, J.C. *see* Sachedina, N.  
 Pieber, T.R. *see* Siebenhofer, A.  
 Pildal, J., Lajer, M., Hansen, S.K., Almind, K., Ambye, L., Borch-Johnsen, K., Carstensen, B., Hansen, T. & Pedersen, O. Studies of variability in the islet amyloid polypeptide gene in relation to Type 2 diabetes, 491  
 Pinach, S. *see* Bo, S.  
 Piolot, R. *see* Mankovsky, B.N.  
 Pirie, F.J. *see* Motala, A.A.  
 Plank, J. *see* Siebenhofer, A.  
 Plaschke, A. *see* Holstein, A.  
 van der Ploeg, H.M. *see* Adriaanse, M.C.  
 Polonsky, W.H. *see* Welch, G.  
 Potter, J. *see* Spruce, M.C.  
 Poulsen, P.L. *see* Hansen, K.W.  
 Poulter, C. *see* Valabhji, J.  
 Pouwels, E. *see* Willemssen, M.A.A.P.  
 Pouwels, M.-J.J., Tack, C.J., Hermus, A.R. & Lutterman, J.A. Treatment with intravenous insulin followed by continuous subcutaneous insulin infusion improves glycaemic control in severely resistant Type 2 diabetic patients, 76  
 Pozzilli, P., Manfrini, S. & Picardi, A. Cow's milk and trials for prevention of Type 1 diabetes, 871  
 Prakash, P.K., Biswas, M., ElBouri, K., Braithwaite, P.A. & Hanna, F.W. Pneumococcal necrotizing fasciitis in a patient with Type 2 diabetes, 899  
 Prasad, Y.D.M. *see* Dang, C.N.  
 Pratas, S. *see* Macedo, A.  
 Price, D. *see* Rance, J.  
 Price, P.A. *see* Masson, E.A.  
 Prieto, J. *see* Botas, P.  
 Propper, C. *see* Bachmann, M.O.  
 Pu, Y.B. *see* Ko, G.T.C.  
 Qiao, Q., Lindström, J., Valle, T.T. & Tuomilehto, J. Progression to clinically diagnosed and treated diabetes from impaired glucose tolerance and impaired fasting glycaemia, 1027  
 Qin, Y.-W. *see* Zhang, J.-L.  
 Qiu, J.-L. *see* Zhang, J.-L.  
 Quentel, G. *see* Massin, P.  
 Rachoń, D., Myśliwska, J., Suchecka-Rachoń, K., Semetkowska-Jurkiewicz, B., Zorena, K. & Kysiak-Szydłowska, W. Serum interleukin-6 levels and bone mineral density at the femoral neck in post-menopausal women with Type 1 diabetes, 475  
 Rahman, M., Griffin, S.J., Rathmann, W. & Wareham, N.J. How should peripheral neuropathy be assessed in people with diabetes in primary care? A population-based comparison of four measures, 368  
 Railton, A. *see* Kaushal, K.  
 Rainbow, S. *see* Sharp, P.S.  
 Ramachandran, A., Snehalatha, C., Satyavani, K. & Vijay, V. Impaired fasting glucose and impaired glucose tolerance in urban population in India, 220  
 Raml, A. *see* Biesenbach, G.  
 Rance, J., Phillips, C., Davies, S., O'Malley, B., Zaman, Q. & Price, D. How much of a priority is treating erectile dysfunction? A study of patients' perceptions, 205  
 Rathmann, W. *see* Rahman, M.  
 Ravikumar, R. *see* Velmurugan, K.  
 Razvi, S., Myers, L., Patton, K. & McCulloch, A.J. Screening for diabetic retinopathy: a cause for concern in people who drive, 812  
 Reid, H. *see* Bingham, E.  
 Renders, C.M., Valk, G.D., de Sonnaville, J.J.J., Twisk, J., Kriegsman, D.M.W., Heine, R.J., van Eijk, J.Th.M. & van der Wal, G. Quality of care for patients with Type 2 diabetes mellitus—a long-term comparison of two quality improvement programmes in the Netherlands, 846  
 Rihl, J. *see* Hattori, N.  
 Roach, P., Woodworth, J., Gudat, U., Cerimele, B., Diebler, F., Pein, M. & Dreyer, M. A 75% insulin lispro/25% NPL mixture provides a longer duration of insulin activity compared with insulin lispro alone in patients with Type 1 diabetes, 946  
 Roberts, C. *see* Vyas, A.  
 Robertson, L. *see* Barnett, A.H.  
 Robinson, R.L. *see* Mazze, R.S.  
 Rodins, K. *see* Tang, E.W.  
 Roland, J. *see* Owen, K.R.  
 Roper, N.A. *see* Vaidya, B.  
 Rosenthal, M. *see* Baldeweg, S.E.  
 Rossi, A.C. *see* Greco, P.  
 Rotella, C.M. *see* Mannucci, E.  
 Rotella, F. *see* Mannucci, E.  
 Rowley, M.J. *see* Davis, T.M.E.  
 Royle, P. *see* Colquitt, J.  
 Rutten, G.E.H.M. *see* Goudswaard, A.N.  
 Rutter, M.K. *see* Fischbacher, C.M.  
 Ruxer, J. *see* Czupryniak, L.  
 Rydén-Ahlgren, Å. *see* Gottsäter, A.  
 Ryder, R.E.J. *see* Basu, A.  
 Saavedra-Gonzalez, Á. *see* García-Estévez, D.A.  
 Saeed, B.O., Banerjee, K., Nixon, S.J. & Brown, K. Plasma homocysteine concentrations in patients with Type 1 diabetes, 867  
 Sachedina, N. & Pickup, J.C. Performance assessment of the Medtronic-MiniMed Continuous Glucose Monitoring System and its use for measurement of glycaemic control in Type 1 diabetic subjects, 1012  
 Salkovskis, P.M. *see* Farmer, A.J.  
 Sandbæk, A. *see* Borch-Johnsen, K.  
 Sandbæk, A. *see* Colagiuri, S.  
 Sanderson, E. *see* Zhao, H.X.  
 Saris, W.H.M. *see* Mensink, M.  
 Särnblad, S. *see* Ingberg, C.-M.

- Saryusz-Wolska, M. *see* Czupryniak, L.  
 Saryavani, K. *see* Ramachandran, A.  
 Sauve, R. *see* Wood, S.L.  
 Scanlon, P. *see* Harding, S.  
 Scanlon, P.H., Malhotra, R., Thomas, G., Foy, C., Kirkpatrick, J.N., Lewis-Barned, N., Harney, B. & Aldington, S.J. The effectiveness of screening for diabetic retinopathy by digital imaging photography and technician ophthalmoscopy, 467  
 Schaefer-Graf, U.M., Kjos, S.L., Bühling, K.J., Henrich, W., Brauer, M., Heinze, T., Dudenhausen, J.W. & Vetter, K. Amniotic fluid insulin levels and fetal abdominal circumference at time of amniocentesis in pregnancies with diabetes, 349  
 Schatz, H. *see* Zimny, S.  
 Scheffer, P.G., Bos, G., Volwater, H.G.F.M., Dekker, J.M., Heine, R.J. & Teerlink, T. Associations of LDL size with *in vitro* oxidizability and plasma levels of *in vivo* oxidized LDL in Type 2 diabetic patients, 563  
 Schmekal, B. *see* Biesenbach, G.  
 Schmidt, K. & Ernst, E. Complementary/alternative medicine for diabetes. How good is advice offered on websites? 248  
 Schnedl, W.J. *see* Lahousen, T.  
 Schnell, O. *see* Hattori, N.  
 Schramm, W., Bergrem, H., Cromme, P., Feest, T., Borch-Johnsen, K., Feldt-Rasmussen, B. & Landgraf, R. First referral of diabetic patients to a nephrologist, 689  
 Schroeder, M. *see* Kaila, B.  
 Schvarcz, E. *see* Ingberg, C-M.  
 Schwaiger, M. *see* Hattori, N.  
 Scioscia, M. *see* Greco, P.  
 Scott, A.R. *see* Dhindsa, P.  
 Scott, R.S. *see* Florkowski, C.M.  
 Scuffham, P. & Carr, L. The cost-effectiveness of continuous subcutaneous insulin infusion compared with multiple daily injections for the management of diabetes, 586  
 Selvaggi, L. *see* Greco, P.  
 Semetkowska-Jurkiewicz, B. *see* Rachoń, D.  
 Senthilvelmurugan, M. *see* Velmurugan, K.  
 Shah, A.M. *see* Wheatcroft, S.B.  
 Shannon, W. *see* Smith, S.M.  
 Sharp, P.F. *see* Olson, J.A.  
 Sharp, P.S., Rainbow, S. & Mukherjee, S. Serum levels of low molecular weight advanced glycation end products in diabetic subjects, 575  
 Shaw, J. *see* Zimmer, P.  
 Shaw, J.E. *see* Tapp, R.J.  
 Shaw, J.E. *see* Williams, J.W.  
 Shiell, A.W. *see* Newsome, C.A.  
 Shier, R. *see* Newsome, C.A.  
 Shin, D.H. *see* Choi, K.M.  
 Shin, J. *see* McFarlane, S.I.  
 Shore, A.C. *see* Konopatskaya, O.  
 Siebenhofer, A., Ng, L.L., Plank, J., Berghold, A., Hödl, R. & Pieber, T.R. Plasma N-terminal pro-brain natriuretic peptide in Type 1 diabetic patients with and without diabetic nephropathy, 535  
 Signorile, A. *see* Bo, S.  
 Simonson, G.D. *see* Mazze, R.S.  
 Simpson, H.L. *see* Holt, R.I.G.  
 Singh, D.L. & Bhattarai, M.D. High prevalence of diabetes and impaired fasting glycaemia in urban Nepal, 170  
 Sinha, B., Brydon, P., Taylor, R.S., Hollins, A., Munro, A., Jenkins, D. & Dunne, F. Maternal ante-natal parameters as predictors of persistent postnatal glucose intolerance: a comparative study between Afro-Caribbeans, Asians and Caucasians, 382  
 Sinha, S. *see* Misra, A.  
 Skiba, M. *see* Krum, H.  
 Skinner, T.C., Howells, L., Greene, S., Edgar, K., McEvilly, A. & Johansson, A. Development, reliability and validity of the Diabetes Illness Representations Questionnaire: four studies with adolescents, 283  
 Skinner, T.C. *see* Parkin, T.  
 Skjærbæk, C. *see* Frystyk, J.  
 Skovlund, S.E. *see* Bott, U.  
 Slattery, J. *see* Morris, A.  
 Smids, D. *see* Cameron, F.J.  
 Smith, K. *see* Dunne, F.  
 Smith, K. *see* Owen, K.R.  
 Smith, S.M., Holohan, J., McAuliffe, A. & Firth, R.G. Irish diabetes detection programme in general practice, 717  
 Smith, S.M., O'Leary, M., Bury, G., Shannon, W., Tynan, A., Staines, A. & Thompson, C. A qualitative investigation of the views and health beliefs of patients with Type 2 diabetes following the introduction of a diabetes shared care service, 853  
 Smith, V. *see* Channon, S.  
 Smith-Laitton, E. *see* Trembl, H.  
 Smyth, C. *see* Bain, S.C.  
 Snehaltha, C. *see* Ramachandran, A.  
 Snoek, F.J. *see* Adriaanse, M.C.  
 So, W.-Y. *see* Ko, G.T.C.  
 Sommerfield, A.J., Ewing, F.M.E., Strachan, M.W.J., Deary, I.J., Aitken, G. & Frier, B.M. Self-treatment of mild symptomatic hypoglycaemia by people with insulin-treated diabetes, 686  
 de Sonnaville, J.J.J. *see* Renders, C.M.  
 Song, Y.D. *see* Kim, D.J.  
 Sönksen, P.H. *see* Holt, R.I.G.  
 Soorani-Lunsing, R.J. *see* Willemsen, M.A.A.P.  
 Soper, C. *see* Zhao, H.X.  
 Sørensen, B. *see* Jensen, D.M.  
 Spijkerman, A.M.W. *see* Adriaanse, M.C.  
 Spruce, M.C., Potter, J. & Coppini, D.V. The pathogenesis and management of painful diabetic neuropathy: a review, 88  
 Staehr, P., Hojlund, K., Hother-Nielsen, O., Holst, J.J. & Beck-Nielsen, H. Does overnight normalization of plasma glucose by insulin infusion affect assessment of glucose metabolism in Type 2 diabetes?, 816  
 Staines, A. *see* Smith, S.M.  
 Standl, E. *see* Hattori, N.  
 Stavrou, P. *see* Basu, A.  
 Steel, J.M. St Vincent's Declaration 10 years on: outcome of diabetic pregnancies, 82  
 Stenhouse, E. *see* Zhao, H.X.  
 Stephenson, C.R. *see* Feltbower, R.G.  
 Stolk, R.P. *see* Goudswaard, A.N.  
 Strachan, M.W.J., Gough, K., McKnight, J.A. & Padfield, P.L. Reply, 509  
 Strachan, F.M. *see* Olson, J.A.  
 Strachan, M.W.J. *see* Sommerfield, A.J.  
 Strachan, M.W.J., Frier, B.M. & Deary, I.J. Type 2 diabetes and cognitive impairment, 1  
 Sturrock, N.D.C. *see* Amin, P.

- Suchecka-Rachon, K. *see* Rachon, D.
- Sugiyama, Y. *see* Nagai, Y.
- Sundkvist, G. *see* Gottsater, A.
- Szelag, B. *see* Gottsater, A.
- Taback, S.P. *see* Kaila, B.
- Tack, C.J. *see* Pouwels, M.-J.J.
- Takemura, Y. *see* Aso, Y.
- Tallon, D. *see* Bachmann, M.O.
- Tan, K.C.B. Feline diabetes—personal experience, 80
- Tang, E.W., Jardine, D.L., Rodins, K. & Evans, J. Respiratory failure secondary to diabetic neuropathy affecting the phrenic nerve, 599
- Tapp, R.J., Shaw, J.E., de Courten, M.P., Dunstan, D.W., Welborn, T.A. & Zimmet, P.Z. Foot complications in Type 2 diabetes: an Australian population-based study, 105
- Tayama, K. *see* Aso, Y.
- Taylor, M.D., Frier, B.M., Gold, A.E. & Deary, I.J. Psychosocial factors and diabetes-related outcomes following diagnosis of Type 1 diabetes in adults: The Edinburgh Prospective Diabetes Study, 135
- Taylor, R. *see* Arun, C.S.
- Taylor, R. *see* Harding, S.
- Taylor, R.S. *see* Sinha, B.
- Taylor, V. *see* Viner, R.M.
- Teerlink, T. *see* Scheffer, P.G.
- Thomas, G. *see* Scanlon, P.H.
- Thomas, G.N., Critchley, J.A.J.H., Tomlinson, B., Cockram, C.S. & Chan, J.C.N. Peripheral vascular disease in Type 2 diabetic Chinese patients: associations with metabolic indices, concomitant vascular disease and genetic factors, 988
- Thomas, N. *see* Ko, G.T.C.
- Thompson, A. *see* Trembl, H.
- Thompson, C. *see* Smith, S.M.
- Thompson, C.J. *see* Moore, K.B.
- Thornton, S.J. *see* Mackie, A.D.R.
- Thorsby, P. *see* Birkeland, K.I.
- Tieszen, K.L. & New, J.P. Alternate site blood glucose testing: do patients prefer it?, 325
- Tomlinson, B. *see* Thomas, G.N.
- Tooke, J.E. *see* Konopatskaya, O.
- Torjesen, P.A. *see* Birkeland, K.I.
- Tormey, W.P. *see* Moore, K.B.
- Trelfa, A. *see* Barnett, A.H.
- Trembl, H., Thompson, A., Smith-Laitton, E., Delaney, P. & Feher, M.D. Striving for equality with insulin delivery devices: not to be 'left' out, 248
- Tsaloumas, M.D. *see* Chatterjee, S.
- Tung, T.-H. *see* Liu, W.-J.
- Tuomilehto, J. *see* Qiao, Q.
- Tuomilehto, J. *see* Williams, J.W.
- Twisk, J. *see* Renders, C.M.
- Tynan, A. *see* Smith, S.M.
- Unwin, N. *see* Parker, L.
- Unwin, N.C. *see* Fischbacher, C.M.
- Vaaler, S. *see* Birkeland, K.I.
- Vaidya, B., Roper, N.A., Connolly, V.M. & Kelly, W.F. Socio-economic deprivation and diabetic foot ulcers: no strong association, 249
- Valabhji, J., Watson, M., Cox, J., Poulter, C., Elwig, C. & Elkeles, R.S. Type 2 diabetes presenting as diabetic ketoacidosis in adolescence, 416
- Valk, G.D. *see* Renders, C.M.
- Valle, T.T. *see* Qiao, Q.
- Velmurugan, K., Deepa, R., Ravikumar, R., Lawrence, J.B., Anshoo, H., Senthilvelmurugan, M., Enas, E.A. & Mohan, V. Relationship of lipoprotein(a) with intimal medial thickness of the carotid artery in Type 2 diabetic patients in south India, 455
- Vetter, K. *see* Schaefer-Graf, U.M.
- Vicaut, E. *see* Massin, P.
- Vice, P.A. *see* Massin, E.A.
- Victor, Z. *see* Massin, P.
- Vijay, V. *see* Ramachandran, A.
- Vimercati, A. *see* Greco, P.
- Viner, R.M., Christie, D., Taylor, V. & Hey, S. Motivational/solution-focused intervention improves HbA<sub>1c</sub> in adolescents with Type 1 diabetes: a pilot study, 739
- Volwater, H.G.F.M. *see* Scheffer, P.G.
- Vora, J.P. *see* Younis, N.
- Vyas, A., Haidery, A.Z., Wiles, P.G., Gill, S., Roberts, C. & Cruickshank, J.K. A pilot randomized trial in primary care to investigate and improve knowledge, awareness and self-management among South Asians with diabetes in Manchester, 1022
- Wake, M. *see* Cameron, F.J.
- van der Wal, G. *see* Renders, C.M.
- Walker, J.D. *see* Massin, E.A.
- Wang, M. *see* Chiu, K.C.
- Warburton, T. Current status of screening for diabetic retinopathy in the UK, 690
- Wareham, N.J. *see* Rahman, M.
- Wareham, N.J. *see* Yuyun, M.F.
- Watkins, P.J., Buxton-Thomas, M.S. & Howard, E.R. Long-term outcome after gastrectomy for intractable diabetic gastroparesis, 58
- Watson, M. *see* Valabhji, J.
- Watts, A.P. *see* Ford-Adams, M.E.
- Waugh, N. *see* Colquitt, J.
- Weinger, K. *see* Welch, G.
- Welborn, T.A. *see* Tapp, R.J.
- Welch, G., Weinger, K., Anderson, B. & Polonsky, W.H. Responsiveness of the Problem Areas In Diabetes (PAID) questionnaire, 69
- Wenby, R.B. *see* Chong-Martinez, B.
- Westergaard, J.G. *see* Jensen, D.M.
- Whatmore, J.L. *see* Konopatskaya, O.
- Wheatcroft, S.B., Williams, I.L., Shah, A.M. & Kearney, M.T. Pathophysiological implications of insulin resistance on vascular endothelial function, 255
- White, M. *see* Fischbacher, C.M.
- White, M. *see* Parker, L.
- Whitford, D.L., Lamont, S.S. & Crosland, A. Screening for Type 2 diabetes: is it worthwhile? Views of general practitioners and practice nurses, 155
- Wiklund, U. *see* Lindmark, S.
- Wild, S. *see* Williams, R.
- Wiles, P.G. *see* Vyas, A.

- Wilkinson-Berka, J. *see* Gilbert, R.E.
- Willemsen, M.A.A.P., Soorani-Lunsing, R.J., Pouwels, E. & Klepper, J. Neuroglycopenia in normoglycaemic patients, and the potential benefit of ketosis, 481
- Williams, R. *see* McFarlane, S.I.
- Williams, I.L. *see* Wheatcroft, S.B.
- Williams, J.W., Zimmet, P.Z., Shaw, J.E., de Courten, M.P., Cameron, A.J., Chitson, P., Tuomilehto, J. & Alberti, K.G.M.M. Gender differences in the prevalence of impaired fasting glycaemia and impaired glucose tolerance in Mauritius. Does sex matter?, 915
- Williams, R. *see* Liu, W.-J.
- Williams, R. & Wild, S. Diabetes prevalence data for the United Kingdom—what do we have and what do we need? 505
- Williams, S. *see* Bachmann, M.O.
- Willis, J.A. *see* Lunt, H.
- Wilson, E. *see* Barton, D.M.
- Winocour, P.H. & Fisher, M. Prediction of cardiovascular risk in people with diabetes, 515
- Wolffenbuttel, B.H.R. *see* Leurs, P.B.
- Wood, E. *see* Yuyun, M.F.
- Wood, S.L., Jick, H. & Sauve, R. The risk of stillbirth in pregnancies before and after the onset of diabetes, 703
- Woodworth, J. *see* Roach, P.
- Working Party of the International Diabetes Federation (European Region). Hypertension in people with Type 2 diabetes: knowledge-based diabetes-specific guidelines, 972
- Worth, R.C. *see* Hassanein, M.M.
- Wredling, R. *see* Bragd, J.
- Wroblewski, M. *see* Gottsäter, A.
- Yamada, K. *see* Pang, H.
- Yamagishi, S.-I. *see* Pang, H.
- Yang, X., Hsu-Hage, B.H.-H., Dong, L., Zhang, H., Zhang, C. & Zhang, Y. Postpartum glucose intolerance in Chinese women with gestational diabetes, 687
- Yeh, W.-T. *see* Pan, W.-H.
- Yen, M.-F. *see* Liu, W.-J.
- Yogev, Y., Ben-Haroush, A., Chen, R., Kaplan, B., Phillip, M. & Hod, M. Continuous glucose monitoring for treatment adjustment in diabetic pregnancies—a pilot study, 558
- Yokoyama, H., Hirasawa, K., Aoki, T., Ishiyama, M. & Koyama, K. Brachial-ankle pulse wave velocity measured automatically by oscillometric method is elevated in diabetic patients with incipient nephropathy, 942
- Yoon, K. *see* Manning, P.J.
- Yoshida, N. *see* Aso, Y.
- Young, R.J. *see* Kaushal, K.
- Younis, N., Broadbent, D.M., Harding, S.P. & Vora, J.P. Incidence of sight-threatening retinopathy in Type 1 diabetes in a systematic screening programme, 758
- Yun, Y.S. *see* Kim, D.J.
- Yuyun, M.F., Dinneen, S.F., Edwards, O.M., Wood, E. & Wareham, N.J. Absolute level and rate of change of albuminuria over 1 year independently predict mortality and cardiovascular events in patients with diabetic nephropathy, 277
- Zaman, Q. *see* Rance, J.
- Zhang, C. *see* Yang, X.
- Zhang, H. *see* Yang, X.
- Zhang, J.-L., Qin, Y.-W., Zheng, X., Qiu, J.-L. & Zou, D.-J. Serum resistin level in essential hypertension patients with different glucose tolerance, 828
- Zhang, L., Krzentowski, G., Albert, A. & Lefebvre, P.J. Factors predictive of nephropathy in DCCT Type 1 diabetic patients with good or poor metabolic control, 580
- Zhang, Y. *see* Yang, X.
- Zhao, H.X., Stenhouse, E., Sanderson, E., Soper, C., Hughes, P., Cross, D., Demaine, A.G. & Millward, B.A. Continued rising trend of childhood Type 1 diabetes mellitus in Devon and Cornwall, England, 168
- Zheng, X. *see* Zhang, J.-L.
- Ziegler, D. *see* Mankovsky, B.N.
- Zimmet, P., Shaw, J. & Alberti, K.G.M.M. Preventing Type 2 diabetes and the dysmetabolic syndrome in the real world: a realistic view, 693
- Zimmet, P.Z. *see* Tapp, R.J.
- Zimmet, P.Z. *see* Williams, J.W.
- Zimny, S., Schatz, H. & Pföhl, U. The effects of applied felted foam on wound healing and healing times in the therapy of neuropathic diabetic foot ulcers, 622
- Zinman, B. *see* Owens, D.R.
- Zorena, K. *see* Rachoń, D.
- Zou, D.-J. *see* Zhang, J.-L.

## Subject Index

- 2-h oral glucose tolerance test, 904
- ACE inhibitor, 708
- adolescence, 416
- adolescent, 394, 739
- adolescents, 1005
- adults, 135
- advanced glycation end products, 575
- aetiology, 88
- aggregation, 301
- albumin excretion rate, 580
- albuminuria, 31, 277, 942
- alternate site glucose testing, 325
- alveolar bone loss, 746
- amniotic fluid insulin, 349
- amputation, 329, 665
- amylin, 491
- anaemia, 661
- analogue insulin, 863
- angiotensin II, 607
- angiotensin-converting enzyme, 677, 988
- antenatal corticosteroids, 73
- anti-hypertensive drugs, 290
- anti-islet cell antibodies, 64
- area under ROC curve, 1027
- Asian Indians, 220, 361
- association, 486
- atherosclerosis, 16, 255, 563, 942
- atorvastatin, 743
- atrial natriuretic peptide, 713
- attitude to health, 155
- attitudes, 152
- autoantibodies, 198
- autoantigen, 198
- automated, 528
- autonomic nervous system, 375, 399

- autonomic neuropathy, 495
- autonomy, 909
- $\beta$ -cell function, 225
- basal insulin, 873
- BASC, 646
- beliefs, 152
- birth weight, 162, 339
- blindness, 186
- blood glucose, 552, 828, 953
- blood glucose meter, 325
- blood glucose self-monitoring, 1012
- blood pressure, 515, 752
- bodycomposition, 1005
- body fat distribution, 552
- body mass index, 23, 182
- brain natriuretic peptide, 535
- buccal, 886
- C reactive protein, 743
- calcium stimulation test, 772
- capillary, 953
- cardiovascular disease, 277, 355, 515, 723, 786
- cardiovascular risk factors, 846
- carotid atherosclerosis, 455, 495, 670
- carotid ultrasonography, 670
- ceftazidime, 242
- central fat, 294
- cerebral autoregulation, 119
- cerebral blood flow velocity, 119
- childhood Type 2 diabetes, 483
- children, 568, 646, 656
- Chinese, 988
- CHQ, 646
- chromosomal translocation, 483
- classification, 329
- clinic, 394
- co-morbidity, 127
- coagulation, 930
- cognitive function, 231
- cohort study, 191
- colour slide, 528
- combination therapy, 935
- communication, 909
- comparability, 953
- complications, 665, 921
- congenital malformations, 734
- continuous glucose monitoring, 558
- control, 921
- coping, 135
- coronary artery disease, 99, 147
- coronary heart disease, 670, 723, 832
- cross-sectional studies, 31
- CSII, 76, 307, 586, 863
- cystic fibrosis, 425
- cystic fibrosis-related diabetes, 425
- D dimer, 930
- DCCT, 580
- Denmark, 307
- dexamethasone, 73
- diabetes, 46, 99, 127, 182, 205, 238, 290, 329, 349, 355, 375, 475, 515, 558, 568, 665, 703, 786, 915, 996, 1022
- diabetes in pregnancy, 858
- diabetes mellitus, 69, 114, 159, 277, 301, 394, 563, 575, 670, 708, 713, 832, 899
- diabetes-related complications, 442
- diabetes screening, 717
- diabetes shared care, 853
- diabetic autonomic neuropathy, 58
- diabetic cardiovascular autonomic neuropathy, 119
- diabetic complications, 808
- diabetic foot, 622, 665
- diabetic foot complications, 105
- diabetic ketoacidosis, 242, 416
- diabetic nephropathy, 661, 677, 808
- diabetic neuropathy, 114, 599
- diabetic retinopathy, 186, 269, 467, 528, 635, 727, 962, 965
- diagnosis, 594
- diagnostic criteria, 51, 904
- diazoxide, 231
- diet, 786
- digital, 528
- digital imaging, 766
- digital photography, 467
- diuretic, 708
- driving, 812
- duration of diabetes, 216
- dyslipidaemia, 406
- economic evaluation, 586
- education, 921
- emotional distress, 69
- endothelial dysfunction, 16, 255
- endothelial function, 375
- eNOS, 838
- epidemiology, 105, 437, 904
- erectile dysfunction, 205, 290
- erythropoietin, 661
- ethnic, 651
- ethnic groups, 31
- ethnicity, 198, 382, 387, 416
- evaluation, 973
- exercise, 786
- factor analysis, 99
- family practice, 155
- fasting plasma glucose, 220, 594
- felted foam, 622
- fetal abdominal circumference, 349
- fetal growth, 858
- fibrates, 515
- foot ulcer, 665
- foot ulceration, 159
- footwear, 665
- Fremantle Diabetes Study, 198
- fundus photography, 635
- gastrectomy, 58
- gastrointestinal, 886
- gastroparesis, 58
- gender, 915
- general practice, 846, 853
- genetics, 677, 988
- gestational diabetes, 594, 703
- gestational diabetes mellitus, 51, 382
- gestational hyperglycaemia, 64
- glibenclamide, 231
- glucose, 339, 451, 462
  - disposal, 816
  - intolerance, 220
  - metabolism, 3
  - production, 816
  - sensing, 238
  - sensor, 1012
- GLUT1-deficiency syndrome, 481
- glutamic acid decarboxylase autoantibodies, 64
- glycaemia, 294
- glycaemic control, 135, 540, 846, 873, 1012
- glycated haemoglobin, 863
- glycated haemoglobin HbA1c, 312
- glycosylated, 540
- grading, 965
- growth hormone, 3, 269
- growth hormone deficiency, 3
- guidelines, 973
- haemoglobin A, 540
- haemostatic parameters, 16
- hazard ratio, 1027
- HbA1c, 162
- HDL-cholesterol, 808
- healthcare costs, 442
- health care professionals, 152
- heart, 375
- heart rate variability, 399
- hepatocyte nuclear factor-1 $\alpha$ , 225
- HOE 901, 545
- HOMA, 294
- home blood glucose monitoring, 325
- homeostasis model assessment, 361
- homicide, 772
- Humalog, 946
- hyperglycaemic symptoms, 319
- hyperinsulinaemia, 406
- hypertension, 290, 708, 828, 973
- hypoglycaemia, 231, 238, 312, 772, 863, 1012, 1016
- hypoglycaemic agents, 540
- IAPP, 491
- ICA, 198



- ICA12, 198  
ICA512/IA-2, 198  
IDDM, 586  
IGF-I, 269  
IGF-II, 269  
IGF binding proteins, 269  
illness beliefs, 283  
image compression, 766  
impaired awareness of hypoglycaemia, 1016  
impaired fasting glucose, 220, 717, 743, 915  
impaired fasting glycemia, 1027  
impaired glucose tolerance, 220, 552, 717, 746, 915, 1027  
incidence, 23, 186, 437  
incidence of diabetes, 1027  
incipient nephropathy, 942  
income, 921  
indapamide, 708  
indicators, 568  
inequality, 921  
infections, 159  
inflammation, 723, 930  
innate immunity, 723  
insulin, 46, 76, 339, 462, 838, 935, 1016  
analogue, 626  
analogues, 312, 946  
aspart, 626  
glargine, 545  
lispro, 46, 656, 946  
lispro high mix, 946  
pump treatment, 307  
pumps, 586  
requirement, 642  
resistance, 3, 64, 76, 99, 255, 361, 425, 462, 500, 651, 816  
secretion, 37, 294  
sensitivity, 37, 399, 425, 743  
insulin-like growth factor, 3  
insulin-treated diabetes, 642  
insulin-treated diabetes mellitus, 191  
intensified insulin therapy, 626  
interleukin-6, 475, 930  
International Diabetes Federation, 973  
intima-media thickness, 670  
intimal medial thickness, 455  
intramyocellular lipid, 361  
intravenous insulin, 73  
ischaemia, 607  
  
Japanese men, 746  
JPEG, 766  
  
ketogenic diet, 481  
ketosis, 481  
knowledge, 1022  
  
LADA, 37  
LAGB, 127  
  
LDL diameter, 563  
LDL oxidation, 563  
leptin, 832  
lifecourse, 406  
lipids, 515  
lipoprotein(a), 455  
lipoproteins, 563  
lyso-PC, 838  
  
macrosomia, 349  
maladjustment, 646  
Markov model, 727  
mass screening, 155  
maternal morbidity, 734  
maternal-fetal outcome, 51  
Mauritius, 915  
melioidosis, 242  
menopause, 475  
meta-analysis, 863  
metabolic control, 1005  
metabolic syndrome, 99, 406, 832, 988  
metabolism, 127  
methicillin-resistant *Staphylococcus aureus*, 159  
microalbuminuria, 713, 808  
microvascular complications, 16, 425  
minimum data set, 965  
modelling, 437  
monofilament, 368  
mortality, 277, 723  
mortality rate, 191  
motivation, 909  
motivational enhancement, 739  
mutation analysis, 491  
  
nasal, 886  
necrotizing fasciitis, 899  
nephropathies, 607  
nephropathy, 277, 535, 580, 786  
neuroglycopenia, 481  
neuropathy, 105, 368, 622, 665  
nitric oxide, 255, 838  
nocturnal hypoglycaemia, 656  
NPH insulin, 545  
nuclear medicine, 375  
nutrition, 786  
  
obesity, 23, 319, 361, 406, 416, 462, 552  
oestrogens, 475  
ophthalmoscopy, 467  
optometrist, 528  
oral, 886  
oral glucose tolerance test, 594, 717  
orthostatic hypo-tension, 119  
osteoporosis, 475  
outcome of care, 486  
outreach visits, 540  
over-reporting, 752  
overnight insulin infusion, 816  
overt nephropathy, 147  
  
paediatric, 394  
PAI-1, 147  
PAID, 69  
painful neuropathy, 88, 114  
patient centred, 909  
patient satisfaction, 486  
patients, 152  
patients' views, 853  
perinatal mortality, 734  
peripheral vascular disease, 105, 988  
pharmacology, 387  
phrenic nerve, 599  
plasma glucose, 23  
plasminogen activator inhibitor-1, 651  
polymorphism, 451, 491  
post-prandial hyperglycaemia, 1012  
postpartum glucose intolerance, 382  
precision, 752  
pregestational diabetes, 683  
pregnancy, 46, 51, 162, 349, 558, 683, 703, 734  
pregnancy outcome, 46  
preterm delivery, 51, 73  
primary care, 540  
priority, 205  
psychology, 739, 996  
public health, 387  
pulmonary, 886  
pulse wave velocity, 942  
pupillary dilatation, 812  
  
qualitative research, 853  
quality, 568  
quality of care, 846  
quality of life, 135, 312, 586, 626  
questionnaires, 996  
  
randomized control trial, 1022  
randomized controlled trial, 935  
recombinant erythropoietin, 661  
recombinant insulin analogue, 545  
red blood cell, 301  
reliability, 283  
renal disease, 575  
renal impairment, 642  
renin-angiotensin system, 988  
repaglinide, 935  
resistin, 828  
respiratory distress syndrome, 73  
respiratory failure, 599  
responsiveness, 69  
retinal screening, 766, 812  
retinopathy, 607, 683  
review, 175  
rheology, 301  
risk factor, 216  
risk factors, 23, 99, 406, 515  
risk perception, 210, 355  
riskpredictor, 1027  
ROC curve, 1027

- routes of insulin delivery, 886
- screening, 175, 210, 382, 467, 528, 594, 635, 646, 727, 904, 962, 965, 996
- secondary failure, 319
- self-care, 909
- self-management, 1022
- self-monitored blood pressure, 752
- sensitivity and specificity, 467
- septicaemia, 242
- severe hypoglycaemia, 216
- shoes, 665
- short 5th metacarpals, 500
- sialic acid, 723
- sight-threatening, 683
- single nucleotide polymorphism, 225
- Soft-Sense meter, 325
- soft-tissue measurement, 858
- solution-focused therapy, 739
- South African Indians, 23
- South Asians, 1022
- SOX-13, 198
- spectral analysis, 399
- standards, 568
- statins, 515
- stillbirth, 703
- Streptococcus pneumoniae*, 899
- sulphonylurea drugs, 772
- sulphonylureas, 231
- surgery, 127, 329
- survey, 182
- symptoms of hypoglycaemia, 1016
- systematic review, 339
- Taiwan, 182
- the metabolic syndrome, 37
- therapeutics, 387
- therapy, 301
- thiazide, 708
- thrombosis, 651
- time trend, 437
- tissue factor pathway inhibitor, 16
- tissue plasminogen activator, 651
- tPA-PAI-1, 147
- transcranial Doppler sonography, 119
- transcription factor 1, 225
- transforming growth factor-beta, 607
- translocation, 838
- treatment, 76, 88, 205, 973
- treatment satisfaction, 626
- trisomy 10p, 483
- Type 1 diabetes, 135, 147, 162, 198, 216, 307, 312, 425, 437, 535, 656, 808, 873, 946, 1005, 1012
- Type 1 diabetes mellitus, 191
- Type 2 diabetes, 37, 76, 152, 155, 175, 198, 210, 319, 368, 399, 416, 425, 486, 491, 677, 734, 746, 853, 873, 930, 988, 1016
- Type 2 diabetes mellitus, 31, 255, 294, 339, 361, 387, 442, 495, 545, 723, 816, 904, 935, 973
- Type 2 (non-insulin-dependent) diabetes, 225
- Type 2 (non-insulin-dependent) diabetes mellitus, 846
- UKPDS, 198, 442
- unawareness, 216
- under-reporting, 752
- urinary albumin excretion, 31
- urinary albumin excretion rate, 677
- validity, 69, 283, 368
- vascular endothelial growth factor, 607
- venous, 953
- vibration threshold, 368
- vitamin D receptor, 451
- VLCD, 319
- waist-hip ratio, 361
- whole blood glucose, 182
- wound, 329
- wound healing, 622
- X chromosome, 483
- Xq deletion, 483



